



Advice for anti-flea and anti ticks products

How to use anti-flea and anti ticks products?

It is vital that you properly assess the prior to administering the product and that it is applied using the appropriate method. To help, here are a few valuable tips on how to use them safely.

A. Detect the presence of fleas on your pet

- Place a blank piece of paper near the base of the tail (hold it or put it down).
- Rub/shake the fur toward the paper.
- If any small, black specks fall off, wet them and crush them with your finger.
- If the paper turns red, fleas are present.

Do not use on cats

B. Follow the basic rules

- Always ask one of our Mondou employees for advice when you buy products.
- Always stay around for several hours following the first use.
- Never use two anti-flea products at the same time.
- If you have more than one pet, separate them for 24 hours after application to prevent them from licking and sharing the product. All products are toxic if ingested.
- Do not give your pet a bath or let it outside in the rain for 24 hours after the treatment.
- Treat your pet once a month for four months, since as the eggs (and nymphs) can survive for four months in the environment.
- All products are toxic if ingested.

Do not use on cats

C. Apply in a safe manner

- Make sure that the product corresponds to the weight AND the animal to be treated.
- Remove one applicator tube from package. See «How to Open» section.
- Hold applicator tube in an upright position facing away from and you and your pet's face and eyes. Pull cap off tube.
- Turn the cap around and place other end of cap back on tube.
- Twist cap to break seal, then remove cap from tube.

- Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. Do not get this product in your pet's eyes, or allow your pet to ingest this product. The product is bitter tasting and salivation may occur for a short time if your pet licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for your pet to lick the product. Do not allow the product to run off.
- Discard empty tube as described on the packaging.
- Under normal conditions this product is effective for a month.
- However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. Do not retreat more often than once every seven (7) days. After flea control is attained, return with a monthly retreatment schedule.

**Important: it is essential to choose a specific product for your pet.
Under no circumstances should the dog product be used on cats.**

D. Watch for potential adverse effects

- Hypersalivation (especially if there is contact with the mouth)
- Lethargy
- Vomiting and/or diarrhea
- Convulsions
- Disorientation and/or loss of balance
- Labored breathing
- Spasms
- Allergic reaction at application site (red rash, itchiness and loss of hair)

E. What to do in case of overdose

If one of these adverse effects appears, we recommend that you:

- Contact immediately 1-888-noflea6 (1-888-663-5326)
- Immediately contact your veterinary clinic.

Your pets' health and well-being are important to us, which is why we are committed to offering you advice on how to choose and properly administer flea prevention and treatment products.